

Guidelines for Teaching Pain Control and Sedation to Dentists and Dental Students

I. INTRODUCTION

The administration of local anesthesia, sedation and general anesthesia is an integral part of the practice of dentistry. The American Dental Association is committed to the safe and effective use of these modalities by appropriately educated and trained dentists.

Anxiety and pain control can be defined as the application of various physical, chemical and psychological modalities to the prevention and treatment of preoperative, operative and postoperative patient anxiety and pain to allow dental treatment to occur in a safe and effective manner. It involves all disciplines of dentistry and, as such, is one of the most important aspects of dental education. The intent of these *Guidelines* is to provide direction for the teaching of pain control and sedation to dentists and can be applied at all levels of dental education from predoctoral through continuing education. They are designed to teach initial competency in pain control and minimal and moderate sedation techniques.

These *Guidelines* recognize that many dentists have acquired a high degree of competency in the use of anxiety and pain control techniques through a combination of instruction and experience. It is assumed that this has enabled these teachers and practitioners to meet the educational criteria described in this document.

It is not the intent of the *Guidelines* to fit every program into the same rigid educational mold. This is neither possible nor desirable. There must always be room for innovation and improvement. They do, however, provide a reasonable measure of program acceptability, applicable to all institutions and agencies engaged in predoctoral and continuing education.

The curriculum in anxiety and pain control is a continuum of educational experiences that will extend over several years of the predoctoral program. It should provide the dental student with the knowledge and skills necessary to provide minimal sedation to alleviate anxiety and control pain without inducing detrimental physiological or psychological side effects. Dental schools whose goal is to have predoctoral students achieve competency in techniques such as local anesthesia and nitrous oxide inhalation and minimal sedation must meet all of the goals, prerequisites, didactic content, clinical experiences, faculty and facilities, as described in these *Guidelines*.

Techniques for the control of anxiety and pain in dentistry should include both psychological and pharmacological modalities. Psychological strategies should include simple relaxation techniques for the anxious patient and more comprehensive behavioral techniques to control pain. Pharmacological strategies should include not only local anesthetics but also sedatives, analgesics and other useful agents. Dentists should learn indications and techniques for administering these drugs enterally, parenterally and by inhalation as supplements to local anesthesia.

The predoctoral curriculum should provide instruction, exposure and/or experience in anxiety and pain control, including minimal and moderate sedation. The predoctoral program must also provide the knowledge and skill to enable students to recognize and manage any emergencies that might arise as a consequence of treatment. Predoctoral

dental students must complete a course in Basic Life Support for the Healthcare Provider. Though Basic Life Support courses are available online, any course taken online should be followed up with a hands-on component and be approved by the American Heart Association or the American Red Cross.

Local anesthesia is the foundation of pain control in dentistry. Although the use of local anesthetics in dentistry has a long record of safety, dentists must be aware of the maximum safe dosage limit for each patient, since large doses of local anesthetics may increase the level of central nervous system depression with sedation. The use of minimal and moderate sedation requires an understanding of local anesthesia and the physiologic and pharmacologic implications of the local anesthetic agents when combined with the sedative agents

The knowledge, skill and clinical experience required for the safe administration of deep sedation and/or general anesthesia are beyond the scope of predoctoral and continuing education programs. Advanced education programs that teach deep sedation and/or general anesthesia to competency have specific teaching requirements described in the Commission on Dental Accreditation requirements for those advanced programs and represent the educational and clinical requirements for teaching deep sedation and/or general anesthesia in dentistry.

The objective of educating dentists to utilize pain control, sedation and general anesthesia is to enhance their ability to provide oral health care. The American Dental Association urges dentists to participate regularly in continuing education update courses in these modalities in order to remain current.

All areas in which local anesthesia and sedation are being used must be properly equipped with suction, physiologic monitoring equipment, a positive pressure oxygen delivery system suitable for the patient being treated and emergency drugs. Protocols for the management of emergencies must be developed and training programs held at frequent intervals.

II. DEFINITIONS

Methods of Anxiety and Pain Control

analgesia — the diminution or elimination of pain.

conscious sedation¹ — a minimally depressed level of consciousness that retains the patient's ability to independently and continuously maintain an airway and respond appropriately to physical stimulation or verbal command and that is produced by a pharmacological or non-pharmacological method or a combination thereof.

In accord with this particular definition, the drugs and/or techniques used should carry a margin of safety wide enough to render unintended loss of consciousness unlikely. Further, patients whose only response is reflex withdrawal from repeated painful stimuli would not be considered to be in a state of conscious sedation.

combination inhalation-enteral conscious sedation (combined conscious sedation) — conscious sedation using inhalation and enteral agents.

When the intent is anxiolysis only, and the appropriate dosage of agents is administered, then the definition of enteral and/or combination inhalation–enteral conscious sedation (combined conscious sedation) does not apply.

local anesthesia — the elimination of sensation, especially pain, in one part of the body by the topical application or regional injection of a drug.

Note: Although the use of local anesthetics is the foundation of pain control in dentistry and has a long record of safety, dentists must always be aware of the maximum, safe dosage limits for each patient. Large doses of local anesthetics in themselves may result in central nervous system depression especially in combination with sedative agents.

minimal sedation — a minimally depressed level of consciousness, produced by a pharmacological method, that retains the patient’s ability to independently and continuously maintain an airway and respond *normally* to tactile stimulation and verbal command. Although cognitive function and coordination may be modestly impaired, ventilatory and cardiovascular functions are unaffected.²

Note: In accord with this particular definition, the drug(s) and/or techniques used should carry a margin of safety wide enough never to render unintended loss of consciousness. Further, patients whose only response is reflex withdrawal from repeated painful stimuli would not be considered to be in a state of minimal sedation.

When the intent is minimal sedation for adults, the appropriate initial dosing of a single enteral drug is no more than the maximum recommended dose (MRD) of a drug that can be prescribed for unmonitored home use.

The use of preoperative sedatives for children (aged 12 and under) prior to arrival in the dental office, except in extraordinary situations, must be avoided due to the risk of unobserved respiratory obstruction during transport by untrained individuals.

Children (aged 12 and under) can become moderately sedated despite the intended level of minimal sedation; should this occur, the guidelines for moderate sedation apply.

For children 12 years of age and under, the American Dental Association supports the use of the American Academy of Pediatrics/American Academy of Pediatric Dentistry *Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures.*

Nitrous oxide/oxygen may be used in combination with a single enteral drug in minimal sedation.

Nitrous oxide/oxygen when used in combination with sedative agent(s) may produce minimal, moderate, deep sedation or general anesthesia.

The following definitions apply to administration of minimal sedation:

maximum recommended dose (MRD) — maximum FDA-recommended dose of a drug as printed in FDA-approved labeling for unmonitored home use.

incremental dosing — administration of multiple doses of a drug until a desired effect is reached, but not to exceed the maximum recommended dose (MRD).

supplemental dosing — during minimal sedation, supplemental dosing is a single additional dose of the initial dose of the initial drug that may be necessary for prolonged procedures. The supplemental dose should not exceed one-half of the initial total dose and should not be administered until the dentist has determined the clinical half-life of the initial dosing has passed. The total aggregate dose must not exceed 1.5x the MRD on the day of treatment.

moderate sedation — a drug-induced depression of consciousness during which patients respond *purposefully* to verbal commands, either alone or accompanied by light tactile stimulation. No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained.³

Note: In accord with this particular definition, the drugs and/or techniques used should carry a margin of safety wide enough to render unintended loss of consciousness unlikely. Repeated dosing of an agent before the effects of previous dosing can be fully appreciated may result in a greater alteration of the state of consciousness than is the intent of the dentist. Further, a patient whose only response is reflex withdrawal from a painful stimulus is not considered to be in a state of moderate sedation.

The following definition applies to administration of moderate and deeper levels of sedation:

titration — administration of incremental doses of a drug until a desired effect is reached. Knowledge of each drug's time of onset, peak response and duration of action is essential to avoid over sedation. Although the concept of titration of a drug to effect is critical for patient safety, when the intent is moderate sedation one must know whether the previous dose has taken full effect before administering an additional drug increment.

deep sedation — a drug-induced depression of consciousness during which patients cannot be easily aroused but respond purposefully following repeated or painful stimulation. The ability to independently maintain ventilatory function may be impaired. Patients may require assistance in maintaining a patent airway, and spontaneous ventilation may be inadequate. Cardiovascular function is usually maintained.³

general anesthesia — a drug-induced loss of consciousness during which patients are not arousable, even by painful stimulation. The ability to independently maintain ventilatory function is often impaired. Patients often require assistance in maintaining a patent airway, and positive pressure ventilation may be required because of depressed spontaneous ventilation or drug-induced depression of neuromuscular function. Cardiovascular function may be impaired.³

3 Excerpted from *Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia*, 2004, of the American Society of Anesthesiologists (ASA). A copy of the full text can be obtained from ASA, 520 N. Northwest Highway, Park Ridge, IL 60068-2573.

Because sedation and general anesthesia are a continuum, it is not always possible to predict how an individual patient will respond. Hence, practitioners intending to produce a given level of sedation should be able to diagnose and manage the physiologic consequences (rescue) for patients whose level of sedation becomes deeper than initially intended.³

For all levels of sedation, the practitioner must have the training, skills, drugs and equipment to identify and manage such an occurrence until either assistance arrives (emergency medical service) or the patient returns to the intended level of sedation without airway or cardiovascular complications.

Routes of Administration

enteral — any technique of administration in which the agent is absorbed through the gastrointestinal (GI) tract or oral mucosa [i.e., oral, rectal, sublingual].

parenteral — a technique of administration in which the drug bypasses the gastrointestinal (GI) tract [i.e., intramuscular (IM), intravenous (IV), intranasal (IN), submucosal (SM), subcutaneous (SC), intraosseous (IO)].

transdermal — a technique of administration in which the drug is administered by patch or iontophoresis through skin.

transmucosal — a technique of administration in which the drug is administered across mucosa such as intranasal, sublingual, or rectal.

inhalation — a technique of administration in which a gaseous or volatile agent is introduced into the lungs and whose primary effect is due to absorption through the gas/blood interface.

Terms

qualified dentist — meets the educational requirements for the appropriate level of sedation in accordance with *Section III* of these *Guidelines*, or a dentist providing sedation and anesthesia in compliance with their state rules and/or regulations prior to adoption of this document.

must/shall — indicates an imperative need and/or duty; an essential or indispensable item; mandatory.

should — indicates the recommended manner to obtain the standard; highly desirable.

may — indicates freedom or liberty to follow a reasonable alternative.

continual — repeated regularly and frequently in a steady succession.

continuous — prolonged without any interruption at any time.

time-oriented anesthesia record — documentation at appropriate time intervals of drugs, doses and physiologic data obtained during patient monitoring.

immediately available — on site in the facility and available for immediate use.

Levels of Knowledge

familiarity — a simplified knowledge for the purpose of orientation and recognition of general principles.

in-depth — a thorough knowledge of concepts and theories for the purpose of critical analysis and the synthesis of more complete understanding (highest level of knowledge).

Levels of Skill

exposed — the level of skill attained by observation of or participation in a particular activity.

competent — displaying special skill or knowledge derived from training and experience.

proficient — the level of skill attained when a particular activity is accomplished with repeated quality and a more efficient utilization of time (highest level of skill).

American Society of Anesthesiologists (ASA) Patient Physical Status Classification⁴

ASA I — A normal healthy patient.

ASA II — A patient with mild systemic disease.

ASA III — A patient with severe systemic disease.

ASA IV — A patient with severe systemic disease that is a constant threat to life.

ASA V — A moribund patient who is not expected to survive without the operation.

ASA VI — A declared brain-dead patient whose organs are being removed for donor purposes.

E — Emergency operation of any variety (used to modify one of the above classifications, i.e., ASA III-E).

Education Courses

Education may be offered at different levels (competency, update, survey courses and advanced education programs). A description of these different levels follows:

1. Competency Courses are designed to meet the needs of dentists who wish to become knowledgeable and proficient in the safe and effective administration of local anesthesia, minimal and moderate sedation. They consist of lectures, demonstrations and sufficient clinical participation to assure the faculty that the dentist understands the procedures taught and can safely and effectively apply them so that mastery of the subject is achieved. Faculty must assess and document the dentist's

III. TEACHING
PAIN CONTROL

competency upon successful completion of such training. To maintain competency, periodic update courses must be completed.

2. **Update Courses** are designed for persons with previous training. They are intended to provide a review of the subject and an introduction to recent advances in the field. They should be designed didactically and clinically to meet the specific needs of the participants. Participants must have completed previous competency training (equivalent, at a minimum, to the competency course described in this document) and have current experience to be eligible for enrollment in an update course.
3. **Survey Courses** are designed to provide general information about subjects related to pain control and sedation. Such courses should be didactic and not clinical in nature, since they are not intended to develop clinical competency.
4. **Advanced Education Courses** are a component of an advanced dental education program, accredited by the ADA Commission on Dental Accreditation in accord with the *Accreditation Standards* for advanced dental education programs. These courses are designed to prepare the graduate dentist or postdoctoral student in the most comprehensive manner to be knowledgeable and proficient in the safe and effective administration of minimal, moderate and deep sedation and general anesthesia.

These *Guidelines* present a basic overview of the recommendations for teaching pain control.

- A. General Objectives:** Upon completion of a predoctoral curriculum in pain control the dentist must:
1. Have an in-depth knowledge of those aspects of anatomy, physiology, pharmacology and psychology involved in the use of various anxiety and pain control methods;
 2. Be competent in evaluating the psychological and physical status of the patient, as well as the magnitude of the operative procedure, in order to select the proper regimen;
 3. Be competent in monitoring vital functions;
 4. Be competent in prevention, recognition and management of related complications;
 5. Be familiar with the appropriateness of and the indications for medical consultation or referral;
 6. Be competent in the maintenance of proper records with accurate chart entries recording medical history, physical examination, vital signs, drugs administered and patient response.

B. Pain Control Curriculum Content:

1. Philosophy of anxiety and pain control and patient management, including the nature and purpose of pain
2. Review of physiologic and psychologic aspects of anxiety and pain
3. Review of airway anatomy and physiology
4. Physiologic monitoring

- a. Observation
 - (1) Central nervous system
 - (2) Respiratory system
 - a. Oxygenation
 - b. Ventilation
 - (3) Cardiovascular system
 - b. Monitoring equipment
5. Pharmacologic aspects of anxiety and pain control
- a. Routes of drug administration
 - b. Sedatives and anxiolytics
 - c. Local anesthetics
 - d. Analgesics and antagonists
 - e. Adverse side effects
 - f. Drug interactions
 - g. Drug abuse
6. Control of preoperative and operative anxiety and pain
- a. Patient evaluation
 - (1) Psychological status
 - (2) ASA physical status
 - (3) Type and extent of operative procedure
 - b. Nonpharmacologic methods
 - (1) Psychological and behavioral methods
 - (a) Anxiety management
 - (b) Relaxation techniques
 - (c) Systematic desensitization
 - (2) Interpersonal strategies of patient management
 - (3) Hypnosis
 - (4) Electronic dental anesthesia
 - (5) Acupuncture/Acupressure
 - (6) Other
 - c. Local anesthesia
 - (1) Review of related anatomy, and physiology

- (2) Pharmacology
 - (a) Dosing
 - (b) Toxicity
 - (c) Selection of agents
- (3) Techniques of administration
 - (a) Topical
 - (b) Infiltration (supraperiosteal)
 - (c) Nerve block — maxilla—to include:
 - (i) Posterior superior alveolar
 - (ii) Infraorbital
 - (iii) Nasopalatine
 - (iv) Greater palatine
 - (v) Maxillary (2nd division)
 - (vi) Other blocks
 - (d) Nerve block — mandible—to include:
 - (i) Inferior alveolar–lingual
 - (ii) Mental–incisive
 - (iii) Buccal
 - (iv) Gow–Gates
 - (v) Closed mouth
 - (e) Alternative injections—to include:
 - (i) Periodontal ligament
 - (ii) Intraosseous

d. Prevention, recognition and management of complications and emergencies

C. Sequence of Pain Control Didactic and Clinical Instruction: Beyond the basic didactic instruction in local anesthesia, additional time should be provided for demonstrations and clinical practice of the injection techniques. The teaching of other methods of anxiety and pain control, such as the use of analgesics and enteral, inhalation and parenteral sedation, should be coordinated with a course in pharmacology. By this time the student also will have developed a better understanding of patient evaluation and the problems related to prior patient care. As part of this instruction, the student should be taught the techniques of venipuncture and physiologic monitoring. Time should be included for demonstration of minimal and moderate sedation techniques.

Following didactic instruction in minimal and moderate sedation, the student must receive sufficient clinical experience to demonstrate competency in those techniques **9**

IV. TEACHING
ADMINISTRATION
OF MINIMAL SEDA-
TION

in which the student is to be certified. It is understood that not all institutions may be able to provide instruction to the level of clinical competence in pharmacologic sedation modalities to all students. The amount of clinical experience required to achieve competency will vary according to student ability, teaching methods and the anxiety and pain control modality taught.

Clinical experience in minimal and moderate sedation techniques should be related to various disciplines of dentistry and not solely limited to surgical cases. Typically, such experience will be provided in managing healthy adult patients. The sedative care of pediatric patients and those with special needs requires advanced didactic and clinical training.

Throughout both didactic and clinical instruction in anxiety and pain control, psychological management of the patient should also be stressed. Instruction should emphasize that the need for sedative techniques is directly related to the patient's level of anxiety, cooperation, medical condition and the planned procedures.

D. Faculty: Instruction must be provided by qualified faculty for whom anxiety and pain control are areas of major proficiency, interest and concern.

E. Facilities: Competency courses must be presented where adequate facilities are available for proper patient care, including drugs and equipment for the management of emergencies.

The faculty responsible for curriculum in minimal sedation techniques must be familiar with the ADA Policy Statement: *Guidelines for the Use of Sedation and General Anesthesia by Dentists*, and the Commission on Dental Accreditation's *Accreditation Standards* for dental education programs.

These *Guidelines* present a basic overview of the recommendations for teaching minimal sedation. These include courses in nitrous oxide/oxygen sedation, enteral sedation, and combined inhalation/enteral techniques.

General Objectives: Upon completion of a competency course in minimal sedation, the dentist must be able to:

1. Describe the adult and pediatric anatomy and physiology of the respiratory, cardiovascular and central nervous systems, as they relate to the above techniques.
2. Describe the pharmacological effects of drugs.
3. Describe the methods of obtaining a medical history and conduct an appropriate physical examination.
4. Apply these methods clinically in order to obtain an accurate evaluation.
5. Use this information clinically for ASA classification and risk assessment.
6. Choose the most appropriate technique for the individual patient.
7. Use appropriate physiologic monitoring equipment.
8. Describe the physiologic responses that are consistent with minimal sedation.
9. Understand the sedation/general anesthesia continuum.

Inhalation Sedation (Nitrous Oxide/Oxygen)

A. Inhalation Sedation Course Objectives: Upon completion of a competency course in inhalation sedation techniques, the dentist must be able to:

1. Describe the basic components of inhalation sedation equipment.
2. Discuss the function of each of these components.
3. List and discuss the advantages and disadvantages of inhalation sedation.
4. List and discuss the indications and contraindications of inhalation sedation.
5. List the complications associated with inhalation sedation.
6. Discuss the prevention, recognition and management of these complications.
7. Administer inhalation sedation to patients in a clinical setting in a safe and effective manner.
8. Discuss the abuse potential, occupational hazards and other untoward effects of inhalation agents.

B. Inhalation Sedation Course Content:

1. Historical, philosophical and psychological aspects of anxiety and pain control.
2. Patient evaluation and selection through review of medical history taking, physical diagnosis and psychological considerations.
3. Definitions and descriptions of physiological and psychological aspects of anxiety and pain.
4. Description of the stages of drug-induced central nervous system depression through all levels of consciousness and unconsciousness, with special emphasis on the distinction between the conscious and the unconscious state.
5. Review of pediatric and adult respiratory and circulatory physiology and related anatomy.
6. Pharmacology of agents used in inhalation sedation, including drug interactions and incompatibilities.
7. Indications and contraindications for use of inhalation sedation.
8. Review of dental procedures possible under inhalation sedation.
9. Patient monitoring using observation and monitoring equipment, with particular attention to vital signs and reflexes related to pharmacology of nitrous oxide.
10. Importance of maintaining proper records with accurate chart entries recording medical history, physical examination, vital signs, drugs and doses administered and patient response.
11. Prevention, recognition and management of complications and life-threatening situations.
12. Administration of local anesthesia in conjunction with inhalation sedation techniques.
13. Description and use of inhalation sedation equipment.
14. Introduction to potential health hazards of trace anesthetics and proposed techniques for limiting occupational exposure.
15. Discussion of abuse potential.

C. Inhalation Sedation Course Duration: While length of a course is only one of the many factors to be considered in determining the quality of an educational program, the course should be a minimum of *14 hours*, including a clinical component during which competency in inhalation sedation technique is achieved. The inhalation sedation course most often is completed as a part of the predoctoral dental education program. However, the course may be completed in a postdoctoral continuing education competency course.

D. Participant Evaluation and Documentation of Inhalation Sedation Instruction: Competency courses in inhalation sedation techniques must afford participants with sufficient clinical experience to enable them to achieve competency. This experience must be provided under the supervision of qualified faculty and must be evaluated. The course director must certify the competency of participants upon satisfactory completion of training. Records of the didactic instruction and clinical experience, including the number of patients treated by each participant must be maintained and available.

E. Faculty: The course should be directed by a dentist or physician qualified by experience and training. This individual should have had at least three years of experience, including the individual's formal postdoctoral training in anxiety and pain control. In addition, the participation of highly qualified individuals in related fields, such as anesthesiologists, pharmacologists, internists, and cardiologists and psychologists, should be encouraged.

A participant-faculty ratio of not more than ten-to-one when inhalation sedation is being used allows for adequate supervision during the clinical phase of instruction; a one-to-one ratio is recommended during the early state of participation.

The faculty should provide a mechanism whereby the participant can evaluate the performance of those individuals who present the course material.

F. Facilities: Competency courses must be presented where adequate facilities are available for proper patient care, including drugs and equipment for the management of emergencies.

Enteral and/or Combination Inhalation-enteral Minimal Sedation

A. Enteral and/or Combination Inhalation-enteral Minimal Sedation Course

Objectives: Upon completion of a competency course in enteral and/or combination inhalation-enteral minimal sedation techniques, the dentist must be able to:

1. Describe the basic components of inhalation sedation equipment.
2. Discuss the function of each of these components.
3. List and discuss the advantages and disadvantages of enteral and/or combination inhalation-enteral minimal sedation (combined minimal sedation).
4. List and discuss the indications and contraindications for the use of enteral and/or combination inhalation-enteral minimal sedation (combined minimal sedation).
5. List the complications associated with enteral and/or combination inhalation-enteral minimal sedation (combined minimal sedation).
6. Discuss the prevention, recognition and management of these complications.

7. Administer enteral and/or combination inhalation–enteral minimal sedation (combined minimal sedation) to patients in a clinical setting in a safe and effective manner.
8. Discuss the abuse potential, occupational hazards and other effects of enteral and inhalation agents.
9. Discuss the pharmacology of the enteral and inhalation drugs selected for administration.
10. Discuss the precautions, contraindications and adverse reactions associated with the enteral and inhalation drugs selected.
11. Describe a protocol for management of emergencies in the dental office and list and discuss the emergency drugs and equipment required for management of life-threatening situations.
12. Demonstrate the ability to manage life-threatening emergency situations, including current certification in Basic Life Support for Healthcare Providers.
13. Discuss the pharmacological effects of combined drug therapy, their implications and their management. Nitrous oxide/oxygen when used in combination with sedative agent(s) may produce minimal, moderate, deep sedation or general anesthesia.

B. Enteral and/or Combination Inhalation–Enteral Minimal Sedation Course Content:

1. Historical, philosophical and psychological aspects of anxiety and pain control.
2. Patient evaluation and selection through review of medical history taking, physical diagnosis and psychological profiling.
3. Definitions and descriptions of physiological and psychological aspects of anxiety and pain.
4. Description of the stages of drug-induced central nervous system depression through all levels of consciousness and unconsciousness, with special emphasis on the distinction between the conscious and the unconscious state.
5. Review of pediatric and adult respiratory and circulatory physiology and related anatomy.
6. Pharmacology of agents used in enteral and/or combination inhalation–enteral minimal sedation, including drug interactions and incompatibilities.
7. Indications and contraindications for use of enteral and/or combination inhalation–enteral minimal sedation (combined minimal sedation).
8. Review of dental procedures possible under enteral and/or combination inhalation–enteral minimal sedation).
9. Patient monitoring using observation, monitoring equipment, with particular attention to vital signs and reflexes related to consciousness.
10. Maintaining proper records with accurate chart entries recording medical history, physical examination, informed consent, time-oriented anesthesia record, including the names of all drugs administered including local anesthetics, doses, and monitored physiological parameters.

11. Prevention, recognition and management of complications and life-threatening situations.
12. Administration of local anesthesia in conjunction with enteral and/or combination inhalation-enteral minimal sedation techniques.
13. Description and use of inhalation sedation equipment.
14. Introduction to potential health hazards of trace anesthetics and proposed techniques for limiting occupational exposure.
15. Discussion of abuse potential.

C. Enteral and/or Combination Inhalation-enteral Minimal Sedation Course

Duration: Participants must be able to document current certification in Basic Life Support for Healthcare Providers and have completed a nitrous oxide competency course to be eligible for enrollment in this course. While length of a course is only one of the many factors to be considered in determining the quality of an educational program, the course should include a minimum of *16 hours*, plus clinically-oriented experiences during which competency in enteral and/or combined inhalation-enteral minimal sedation techniques is demonstrated. Clinically-oriented experiences may include group observations on patients undergoing enteral and/or combination inhalation-enteral minimal sedation. Clinical experience in managing a compromised airway is critical to the prevention of life-threatening emergencies. The faculty should schedule participants to return for additional clinical experience if competency has not been achieved in the time allotted. The educational course may be completed in a predoctoral dental education curriculum or a postdoctoral continuing education competency course.

These *Guidelines* are not intended for the management of enteral and/or combination inhalation-enteral minimal sedation in children, which requires additional course content and clinical learning experience.

D. Participant Evaluation and Documentation of Instruction: Competency courses in combination inhalation-enteral minimal sedation techniques must afford participants with sufficient clinical understanding to enable them to achieve competency. The course director must certify the competency of participants upon satisfactory completion of the course. Records of the course instruction must be maintained and available.

E. Faculty: The course should be directed by a dentist or physician qualified by experience and training. This individual should have had at least three years of experience, including the individual's formal postdoctoral training in anxiety and pain control. Dental faculty with broad clinical experience in the particular aspect of the subject under consideration should participate. In addition, the participation of highly qualified individuals in related fields, such as anesthesiologists, pharmacologists, internists, and cardiologists and psychologists, should be encouraged. The faculty should provide a mechanism whereby the participant can evaluate the performance of those individuals who present the course material.

F. Facilities: Competency courses must be presented where adequate facilities are available for proper patient care, including drugs and equipment for the management of emergencies.

V. TEACHING
ADMINISTRATION OF
MODERATE SEDATION

These *Guidelines* present a basic overview of the requirements for a competency course in moderate sedation. These include courses in enteral moderate sedation and parenteral moderate sedation. The teaching guidelines contained in this section on moderate sedation differ slightly from documents in medicine to reflect the differences in delivery methodologies and practice environment in dentistry. For this reason, separate teaching guidelines have been developed for moderate enteral and moderate parenteral sedation.

A. Course Objectives: Upon completion of a course in moderate sedation, the dentist must be able to:

1. List and discuss the advantages and disadvantages of moderate sedation.
2. Discuss the prevention, recognition and management of complications associated with moderate sedation.
3. Administer moderate sedation to patients in a clinical setting in a safe and effective manner.
4. Discuss the abuse potential, occupational hazards and other untoward effects of the agents utilized to achieve moderate sedation.
5. Describe and demonstrate the technique of intravenous access, intramuscular injection and other parenteral techniques.
6. Discuss the pharmacology of the drug(s) selected for administration.
7. Discuss the precautions, indications, contraindications and adverse reactions associated with the drug(s) selected.
8. Administer the selected drug(s) to dental patients in a clinical setting in a safe and effective manner.
9. List the complications associated with techniques of moderate sedation.
10. Describe a protocol for management of emergencies in the dental office and list and discuss the emergency drugs and equipment required for the prevention and management of emergency situations.
11. Discuss principles of advanced cardiac life support or an appropriate dental sedation/anesthesia emergency course equivalent.
12. Demonstrate the ability to manage emergency situations.

B. Moderate Sedation Course Content:

1. Historical, philosophical and psychological aspects of anxiety and pain control.
2. Patient evaluation and selection through review of medical history taking, physical diagnosis and psychological considerations.
3. Definitions and descriptions of physiological and psychological aspects of anxiety and pain.
4. Description of the sedation anesthesia continuum, with special emphasis on the distinction between the conscious and the unconscious state.
5. Review of pediatric and adult respiratory and circulatory physiology and related anatomy.
6. Pharmacology of local anesthetics and agents used in moderate sedation, including drug interactions and contraindications.

7. Indications and contraindications for use of moderate sedation.
8. Review of dental procedures possible under moderate sedation.
9. Patient monitoring using observation and monitoring equipment, with particular attention to vital signs and reflexes related to consciousness.
10. Maintaining proper records with accurate chart entries recording medical history, physical examination, informed consent, time-oriented anesthesia record, including the names of all drugs administered including local anesthetics, doses, and monitored physiological parameters.
11. Prevention, recognition and management of complications and emergencies.
12. Description and use of moderate sedation monitors and equipment.
13. Discussion of abuse potential.
14. Intravenous access: anatomy, equipment and technique.
15. Prevention, recognition and management of complications of venipuncture and other parenteral techniques.
16. Description and rationale for the technique to be employed.
17. Prevention, recognition and management of systemic complications of moderate sedation, with particular attention to airway maintenance and support of the respiratory and cardiovascular systems.

C. Moderate Enteral Sedation Course Duration: A minimum of *24 hours* of instruction, plus management of at least *10 adult case experiences* by the enteral and/or enteral-nitrous oxide/oxygen route are required to achieve competency. These ten cases must include at least three live clinical dental experiences managed by participants in groups no larger than five. The remaining cases may include simulations and/or video presentations, but must include one experience in returning (rescuing) a patient from deep to moderate sedation. Participants combining enteral moderate sedation with nitrous oxide-oxygen must have first completed a nitrous oxide competency course.

Participants should be provided supervised opportunities for clinical experience to demonstrate competence in airway management. Clinical experience will be provided in managing healthy adult patients; **this course in moderate enteral sedation is not designed for the management of children (aged 12 and under)**. Additional supervised clinical experience is necessary to prepare participants to manage medically compromised adults and special needs patients. This course in moderate enteral sedation does not result in competency in moderate parenteral sedation. The faculty should schedule participants to return for additional didactic or clinical exposure if competency has not been achieved in the time allotted.

Moderate Parenteral Sedation Course Duration: A minimum of *60 hours* of instruction, plus management of *at least 20 patients* by the intravenous route per participant, is required to achieve competency in moderate sedation techniques. Participants combining parenteral moderate sedation with nitrous oxide-oxygen must have first completed a nitrous oxide competency course.

Clinical experience in managing a compromised airway is critical to the prevention of emergencies. Participants should be provided supervised opportunities for clinical experience to demonstrate competence in management of the airway. Typically, clinical experience will be provided in managing healthy adult patients.

Additional supervised clinical experience is necessary to prepare participants to manage children (aged 12 and under) and medically compromised adults.

Successful completion of this course does result in clinical competency in moderate parenteral sedation. The faculty should schedule participants to return for additional clinical experience if competency has not been achieved in the time allotted.

D. Participant Evaluation and Documentation of Instruction: Competency courses in moderate sedation techniques must afford participants with sufficient clinical experience to enable them to achieve competency. This experience must be provided under the supervision of qualified faculty and must be evaluated. The course director must certify the competency of participants upon satisfactory completion of training in each moderate sedation technique, including instruction, clinical experience and airway management. Records of the didactic instruction and clinical experience, including the number of patients managed by each participant in each anxiety and pain control modality must be maintained and available for review.

E. Faculty: The course should be directed by a dentist or physician qualified by experience and training. This individual should have had at least three years of experience, including formal postdoctoral training in anxiety and pain control. Dental faculty with broad clinical experience in the particular aspect of the subject under consideration should participate. In addition, the participation of highly qualified individuals in related fields, such as anesthesiologists, pharmacologists, internists, cardiologists and psychologists, should be encouraged.

A participant-faculty ratio of not more than five-to-one when moderate enteral sedation is being taught allows for adequate supervision during the clinical phase of instruction. A participant-faculty ratio of not more than three-to-one when moderate parenteral sedation is being taught allows for adequate supervision during the clinical phase of instruction; a one-to-one ratio is recommended during the early stage of participation.

The faculty should provide a mechanism whereby the participant can evaluate the performance of those individuals who present the course material.

F. Facilities: Competency courses in moderate sedation must be presented where adequate facilities are available for proper patient care, including drugs and equipment for the management of emergencies. These facilities may include dental and medical schools/offices, hospitals and surgical centers.

V. ADDITIONAL SOURCES
OF INFORMATION

American Dental Association. Example of a time oriented anesthesia record at ADA.org.

American Academy of Pediatric Dentistry (AAPD). *Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures: An Update*. Developed through a collaborative effort between the American Academy of Pediatrics and the AAPD. Available at www.aapd.org/policies.

American Academy of Periodontology (AAP). *Guidelines: In-Office Use of Conscious Sedation in Periodontics*. Available at www.perio.org/resources-products/posppr3-1.html. **The AAP rescinded this policy in 2008.**

American Association of Oral and Maxillofacial Surgeons (AAOMS). *Parameters and Pathways: Clinical Practice Guidelines for Oral and Maxillofacial Surgery (AAOMS ParPath o1) Anesthesia in Outpatient Facilities*. Contact AAOMS at 847.678.6200 or visit www.aaoms.org/index.php.

American Association of Oral and Maxillofacial Surgeons (AAOMS). *Office Anesthesia Evaluation Manual 7th Edition*. Contact AAOMS at 847.678.6200 or visit www.aaoms.org/index.php.

American Society of Anesthesiologists (ASA). *Practice Guidelines for Preoperative Fasting and the Use of Pharmacological Agents to Reduce the Risk of Pulmonary Aspiration: Application to Healthy Patients Undergoing Elective Procedures*. Available at <https://ecommerce.asahq.org/p-178-practice-guidelines-for-preoperative-fasting.aspx>.

American Society of Anesthesiologists (ASA). *Practice Guidelines for Sedation and Analgesia by Non-Anesthesiologists*. Available at www.asahq.org/Home/For-Members/Practice-Management/Practice-Parameters#sedation.

The ASA has other anesthesia resources that might be of interest to dentists. For more information, go to www.asahq.org/publicationsAndServices/sgstoc.htm.

Commission on Dental Accreditation (CODA). *Accreditation Standards for Predoctoral and Advanced Dental Education Programs*. Available at ADA.org/115.aspx.

National Institute for Occupational Safety and Health (NIOSH). *Controlling Exposures to Nitrous Oxide During Anesthetic Administration* (NIOSH Alert: 1994 Publication No. 94-100). Available at www.cdc.gov/niosh/docs/94-100/.

Dionne, Raymond A.; Yagiela, John A., et al. Balancing efficacy and safety in the use of oral sedation in dental outpatients. *JADA* 2006;137(4):502-13. ADA members can access this article online at jada.ada.org/cgi/content/full/137/4/502.